

Note

Some upper bounds for the product of the domination number and the chromatic number of a graph

Jerzy Topp* and Lutz Volkmann

Lehrstuhl II für Mathematik, RWTH Aachen, Templergraben 55, 5100 Aachen, Germany

Received 11 December 1991

Abstract

Topp, J. and L. Volkmann, Some upper bounds for the product of the domination number and the chromatic number of a graph, *Discrete Mathematics* 118 (1993) 289–292.

Some new upper bounds for $\gamma\chi$ are proved, where γ is the domination number and χ is the chromatic number of a graph.

All graphs considered in this note are finite, undirected, and without loops or multiple edges. Our terminology is based on [2]. For a graph G , let n , δ , Δ , γ , and χ denote the order, the minimum degree, the maximum degree, the domination number, and the chromatic number of G , respectively. Recently Gernert [5] has obtained the following two inequalities for the product $\gamma\chi$ of the domination number and the chromatic number of a graph.

Theorem A. *If G is a connected graph with $n \geq 5$, then $\gamma\chi \leq \frac{1}{4}n^2$.*

Theorem B. *If G is a regular graph with $n \geq 5$, then $\gamma\chi \leq \frac{6}{25}n^2$.*

In this note we present some improvements of the above inequalities. The following two lemmas will be useful in our proofs.

Correspondence to: Jerzy Topp, Faculty of Applied Physics and Mathematics, Gdańsk Technical University, Majakowskiego 11/12, 80-952 Gdańsk, Poland.

*Research supported by the Heinrich Hertz Foundation.

Lemma 1. For a connected graph G , the following inequalities hold:

- (i) ([1]) $\chi \leq \Delta$ if G is neither a complete graph nor an odd cycle;
- (ii) ([3]) $\chi \leq 3n/5$ if G is a regular graph with $\Delta \leq n-2$;
- (iii) ([6]) $\gamma \leq (n+1-\delta)/2$ if G is not isomorphic to the complement of a one-regular graph;
- (iv) ([6])

$$\gamma \leq 2 + \frac{(n-1-\Delta)(n-2-\delta)}{n-1} \leq \frac{f_{n,\delta}(\Delta)}{\Delta};$$

- (v) ([4])

$$\gamma \leq \frac{n+1}{2} - \frac{(\delta-1)\Delta}{2\delta} \leq \frac{g_{n,\delta}(\Delta)}{\Delta}.$$

Lemma 2. For integers n and δ with $n > \delta + 2 \geq 4$, let $f_{n,\delta}$ and $g_{n,\delta}$ be the functions defined in Lemma 1. Then $f_{n,\delta}$ is decreasing in the interval $(\Delta_{n,\delta}^f, +\infty)$, $g_{n,\delta}$ is increasing in the interval $(-\infty, \Delta_{n,\delta}^g)$, and $g_{n,\delta}$ has the maximum at the point $\Delta_{n,\delta}^g$, where

$$\Delta_{n,\delta}^f = \frac{(n-1)(n-\delta)}{2(n-2-\delta)} \quad \text{and} \quad \Delta_{n,\delta}^g = \frac{\delta(n+1)}{2(\delta-1)}.$$

Theorem 1. If G is a connected regular graph with $n \geq 6$ and G is different from the cycle C_7 , then

$$\gamma\chi \leq \frac{(n+1)^2}{8}. \quad (1)$$

Proof. If G is a complete graph, then $\gamma\chi = n$ and (1) holds for $n \geq 6$. If G is a cycle, then $\gamma\chi = \lceil n/3 \rceil (2-n+2\lceil n/2 \rceil)$ and therefore (1) holds if $n=6$ or $n \geq 8$. Thus we may assume that G is neither a complete graph nor a cycle. Then $3 \leq \Delta \leq n-2$ and we consider two cases. First, if n is odd or $\Delta \neq n-2$, then combining (i) and (iii) we obtain

$$\gamma\chi \leq \frac{(n+1-\Delta)\Delta}{2} \leq \max_{\Delta} \frac{(n+1-\Delta)\Delta}{2} = \frac{(n+1)^2}{8}.$$

Second, assume that $\Delta = n-2$ with even $n \geq 6$. Then $\gamma = 2$ and, according to (ii), $\chi \leq 3n/5$ (resp., $\chi \leq 3$ if $n=6$). Hence,

$$\gamma\chi \leq \frac{6n}{5} \leq \frac{(n+1)^2}{8}$$

for $n \geq 8$, while $\gamma\chi \leq 6 \leq (6+1)^2/8$ for $n=6$. \square

Theorem 2. If G is a connected graph with $\delta \geq 2$, then

$$\gamma\chi \leq \frac{\delta}{8(\delta-1)} (n+1)^2. \quad (2)$$

Proof. It is easy to verify the inequality (2) if G is a complete graph or a cycle. If G is neither a complete graph nor a cycle, then applying (i), (v), and Lemma 2 we obtain

$$\gamma\chi \leq g_{n,\delta}(\Delta) \leq g_{n,\delta}(\Delta_{n,\delta}^g) = \frac{\delta}{8(\delta-1)}(n+1)^2. \quad \square$$

Note that Theorem B follows from Theorem 1 for connected regular graphs of order at least 6 which are different from the cycle C_7 . Similarly, Theorem A follows from Theorem 2 if $\delta \geq 4$ or if $\delta = 3$ and $n \geq 7$. Finally, for $\delta = 2$ and $n \geq 17$, Theorem A follows from the next result.

Theorem 3. *If G is a connected graph with $\delta = 2$, then*

$$\gamma\chi \leq \frac{2}{9}(n+1)^2. \quad (3)$$

Proof. A simple verification shows that the inequality (3) holds if G is a cycle. Thus assume that G is not a cycle. Then the hypothesis $\delta = 2$ implies that G is not a complete graph and $n \geq 4$. There are two cases to consider.

Case 1: $\Delta \leq 2(n+1)/3$.

Since the function $g_{n,2}$ is increasing in $(-\infty, \Delta_{n,2}^g)$ (see Lemma 2) and

$$\Delta \leq 2(n+1)/3 < n+1 = \Delta_{n,2}^g,$$

so we have

$$g_{n,2}(\Delta) \leq g_{n,2}(2(n+1)/3) = 2(n+1)^2/9.$$

Consequently, it follows from (i) and (v) that

$$\gamma\chi \leq g_{n,2}(\Delta) \leq \frac{2}{9}(n+1)^2.$$

Case 2: $\Delta > 2(n+1)/3$.

In this case n must be not smaller than 6. First, if $n = 6$, then $\chi \leq 5$, $\Delta = 5$ and therefore $\gamma = 1$, so $\gamma\chi \leq 5$ and the inequality (3) holds. Second, if $n \geq 7$, then

$$\Delta_{n,2}^f \leq 2(n+1)/3 < \Delta$$

and therefore

$$f_{n,2}(\Delta) \leq f_{n,2}(2(n+1)/3)$$

(by Lemma 2). This together with (i) and (iv) gives

$$\gamma\chi \leq f_{n,2}(\Delta) \leq f_{n,2}(\frac{2}{3}(n+1)) \leq \frac{2}{9}(n+1)^2. \quad \square$$

Concluding remarks. For integers $r \geq 1$ and $k \geq 2$, let G be a graph formed as follows: Take $n = rk + k$ vertices $x_1, x_2, \dots, x_{rk}, a_1, \dots, a_k$ and join x_i to x_j , $1 \leq i < j \leq rk$. Further, join a_i to the vertices $x_{(i-1)r+1}, \dots, x_{ir}$, $1 \leq i \leq k$. In the resulting graph G we have $\delta = r$, $\chi = \delta k$, $\gamma = k$, and

$$\gamma\chi = \delta k^2 = \frac{\delta}{(\delta+1)^2} n^2. \quad (4)$$

For positive integers n and δ with $n > \delta$, let $B_{n,\delta}$ be the smallest integer B such that the inequality $\gamma\chi \leq B$ holds for every connected graph of order n and minimum degree δ . It follows from (4) and Theorem 2 that

$$\frac{\delta}{(\delta+1)^2} n^2 \leq B_{n,\delta} \leq \frac{\delta}{8(\delta-1)} (n+1)^2$$

for $\delta \geq 2$. In particular, for $\delta = 3$ we have

$$\frac{3}{16} n^2 \leq B_{n,3} \leq \frac{3}{16} (n+1)^2.$$

Consequently, the estimation (2) is asymptotically best possible for $\delta = 3$. In a quite similar way, we obtain that the estimation (3) is asymptotically best possible for $\delta = 2$. Finally, let us observe that in the above defined graph G for $k = r$ we have $k = r = \gamma = \delta$, $\Delta = \delta^2$, and (v) yields $\gamma \leq g_{n,\delta}(\delta^2)/\delta^2 = \delta + 1/2$. This implies that the inequality (v) in Lemma 1 is, in some sense, best possible.

References

- [1] R.L. Brooks, On colouring the nodes of a network, *Proc. Cambridge Philos. Soc.* 37 (1941) 194–197.
- [2] G. Chartrand and L. Lesniak, *Graphs and Digraphs* (Wadsworth, Monterey, 1986).
- [3] P. Erdős and T. Gallai, Solution of a problem of Dirac, in: M. Fiedler, ed., *Theory of Graphs and its Applications*, *Proc. Symp. Smolenice 1963* (Academia, Prague, 1964), 167–168.
- [4] P. Flach and L. Volkmann, Estimations for the domination number of a graph, *Discrete Math.* 80 (1990) 145–151.
- [5] D. Gernert, Inequalities between the domination number and the chromatic number of a graph, *Discrete Math.* 76 (1989) 151–153.
- [6] C. Payan, Sur le nombre d'absorption d'un graph simple, *Cahiers Centre Études Rech. Opér.* 17 (1975) 307–317.